

# CHAMOT

## Recombinant Mouse RANKL

CM045-5MP  
CM045-20MP  
CM045-100MP  
CM045-500MP  
CM045-1000MP



CHAMOT

喬默生物技術(上海)有限公司  
CHAMOT BIOTECHNOLOGY CO., LTD.

# CONTENT

1 产品简介

2 产品组成

3 产品储存


4 产品使用

## Recombinant Mouse RANKL (Receptor activator of nuclear factor kappa- B Ligand)

|     |   |     |   |
|-----|---|-----|---|
| 编号: | CM045-5MP<br>CM045-20MP<br>CM045-100MP<br>CM045-500MP<br>CM045-1000MP | 规格: | 5 µg<br>20 µg<br>100 µg<br>500 µg<br>1 mg |
| 类别: | 重组蛋白  | 应用: | Functional Assay                          |

### 产品简介

**描述:** RANKL and RANK are members of the TNF superfamily of ligands and receptors that is critical for the regulation of specific immunity and bone turnover. RANK receptor was originally identified as a dendritic cell -membrane protein, which, by interacting with RANKL, augments the ability of dendritic cells. These dendritic cells then stimulate naïve T-cell proliferation in a mixed lymphocyte reaction, promote the survival of RANK+ T-cells, and regulate T-cell-dependent immune response. RANKL, which is expressed in a variety of cells, including osteoblasts, fibroblasts, activated T-cells and bone marrow stromal cells, is also capable of interacting with a decoy receptor called OPG.



SDS- PAGE analysis of recombinant mouse RANKL

**来源:** *Escherichia coli*

**纯度:** >98% as determined by SDS-PAGE. Ni- NTA chromatography.

**生物学活性:** Measure by its ability to induce osteoclast differentiation in RAW264.7 cells. The ED<sub>50</sub> for this effect is < 2 ng/mL.

**内毒素检测:** < 0.1 EU per 1 µg of the protein by the LAL method.

**蛋白序列:** MPAMMEGSWLDVAQRGKPEAQPF  
 HLTINAASIPSGSHKVTLSWYHDRG

WAKISNMTLSNGKLRVNQDGFYYLY  
ANICFRHHETSGSVPTDYLQLMVYVV  
KTSIKIPSSHNLKGGSTKNWSGNSE  
HFYSINVGFFKLRAGEEISIQVSNPSL  
LDPDQDATYFGAFKVQDID with  
polyhistidine tag at the C-terminus

## 产品组成

成分: 从含有 1X PBS, pH 8.0 溶液中冻干的蛋白质.

## 产品储存/运输

| 产品形式      | 储存温度        | 储存时间     |
|-----------|-------------|----------|
| 冻干粉       | -20°C至-80°C | 自收到之日起1年 |
| 重悬液 (初始)  | 2°C至8°C     | 不超过1周    |
| 重悬液 (经稀释) | -20°C至-80°C | 3到6个月    |

运输方式: 蓝冰

## 产品使用

- 1、开盖前, 建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉, 溶液浓度不低于100 $\mu$ g/mL, 不高于1mg/mL, 并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液, 2-8°C无菌保存不超过1周。
- 4、如需长期保存, 推荐使用无菌的含载体蛋白 (如0.1%BSA、10%FBS 或5%HSA) 的溶液进一步稀释 (不低于10 $\mu$ g/mL) 后分装保存, -20°C至-80°C可无菌保存3到6个月。无血清实验需求时, 可更换为5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;  
FACS= Fluorescence activated Cell Sorting; FA= Functional Assay